

The Structure of EF-Tu-Kirromycin Complex: Implications for the Bacteriostatic and Bacteriocidal Action of the Antibiotic

S. Heffron, A. Chandler, S. Watkins, and F. Journak
Dept. of Physiology and Biophysics, 346-D Med. Sci. I
University of California, Irvine, California 92697-4560, USA

ABSTRACT

In the presence of the antibiotic, kirromycin, *Escherichia coli* elongation factor Tu (EF-Tu) adopts a conformation that remains bound to the ribosome, even after GTP hydrolysis, thus inhibiting further cycles of elongation. The three-dimensional structure of a complex between *E. coli* EF-Tu:GDP and kirromycin has been determined by X-ray diffraction analysis to a resolution of 2.2Å. The overall conformation of the complex is distinct from all previously observed EF-Tu structures, but is most similar to that of EF-Tu-GTP. Not only are the relative domain arrangements different from those observed in other complexes, but the antibiotic also alters the conformation of the Effector Loops I and II. Several kirromycin-resistant mutants have been identified and all sites map to the interface between Domains I and III [1,2]. A number of studies have suggested that the chromophoric end of kirromycin is involved in the interaction with EF-Tu. The present three-dimensional structure of EF-Tu-GDP-kirromycin explains all of the observed biochemical data. In its bacteriostatic mode, kirromycin alters the behavior of EF-Tu such that the activities normally elicited by EF-Ts, aminoacyl-tRNA and the ribosome are mimicked [3]. The new conformation of EF-Tu, in the presence of kirromycin, provides a structural explanation for all observed effects of kirromycin on each of the known EF-Tu complexes. The bacteriocidal mode is less well understood. It may be related to the observation that kirromycin promotes the formation of EF-Tu filaments [4], thereby removing all functional EF-Tu from the cell. An EF-Tu-GDP-kirromycin filament is observed in the crystal which may help to elucidate the molecular basis for the induced filament formation.

REFERENCES

1. F. Abdulkarim, L. Liljas, D. Hughes, "Mutations to kirromycin resistance occur in the interface of domains I and III of EF-Tu-GTP," FEBS Letters **352**, 118-122 (1994).
2. C. Cappellano, F. Monti, M. Sosio, S. Donadio, and E. Sarubbi, "Natural kirromycin resistance of elongation factor Tu from the kirrothricin producer *Streptomyces cinnamomeus*," Microbiology **143**, 617-624 (1997).
3. A. Parmeggiani and G.W.M. Swart, "Mechanism of action of kirromycin-like antibiotics," Ann. Rev. Microbiol. **39**, 557-77 (1985).
4. B.D. Beck, "Polymerization of the bacterial elongation factor for protein synthesis, EF-Tu," Eur. J. Biochem. **97**, 495-502 (1979).

This work was supported by NIH Grant GM26895 to Frances Journak.

Principal investigator: Frances Journak, Dept. of Physiology and Biophysics, College of Medicine, University of California, Irvine. Email: journak@uci.edu. Telephone: 949-824-6623.